

***Pomaderris angustifolia* N.A. Wakef.**

*P. angustifolia* is a multi-branched shrub to 3m tall found in open forest in gullies, often near streams.

**Distribution and Abundance:** A wide-ranging species with disjunct populations from north-eastern NSW and along the Great Dividing Range to near Maffra, Vic. More common in Southern Tablelands of NSW.

**Local Distribution:** It is fairly common in the ACT and surrounding NSW with significant populations along watercourses, including the Murrumbidgee Corridor, Molonglo River, Queanbeyan River as well as in Tidbinbilla Nature Reserve and parts of Namadgi National Park.

**Similar Species:** *P. helianthemifolia* ssp. *hispidula*, from which it differs in having stellate hairs on the upper leaf surface rather than simple hairs.



Kowen Forest



Stems stellate-tomentose



Leaves ± narrow-oblong, 0.7-1.5cm long, about 2mm wide



Leaf margins recurved. Upper surface bristly with scattered stellate hairs, lower surface whitish with dense covering of stellate hairs



Stellate hairs on upper leaf surface



Buds whitish tomentose



Flowers creamish in narrow leafy panicles



Ovary glabrous (green)



Petals absent



Capsules glabrous, capsule almost entirely exserted from shortly tomentose hypanthium